

Interpretation of the Wigner Energy as due to RPA Correlations

Kai Neergård

Næstved Gymnasium og HF, Nygårdsvej 43, DK-4700 Næstved, Denmark,
neergard@inet.uni2.dk

Abstract

In a schematic model with equidistant fourfold degenerate single-nucleon levels, a conventional isovector pairing force and a symmetry force, the RPA correlation energy rises almost linearly with the isospin T , thus producing a Wigner term in accordance with the empirical proportionality of the symmetry energy to $T(T+1)$.

Key words:

symmetry energy, Wigner energy, isovector pairing force, symmetry force, isobaric invariance, RPA, Goldstone mode

PACS-numbers: 21.10.Dr, 21.60.Jz, 27.40.+z

Nearly symmetric nuclei have an extra binding, the so-called Wigner energy, that is not described by the quadratic symmetry term in the semi-empirical mass formula [1]. This is explained in various ways in the literature. Counting the even bonds among supermultiplet degenerate nucleon orbitals, Wigner estimates that the isospin-dependent part of the interaction energy in the ground state of a doubly even nucleus is proportional to $T(T+4)$, where T is the isospin [2]. Talmi proves that for seniority-conserving forces such as the pairing force acting in a single j -shell, this part of the interaction energy is proportional to $T(T+1)$ [3], and Bohr and Mottelson point out that this isospin-dependence arises in general from an interaction proportional to the scalar product of the isospins of the interacting nucleons [4]. A symmetry energy proportional to $T(T+1)$ also fits the empirical masses well [5]. Myers and Swiatecki attribute the extra binding of nearly symmetric nuclei to the interaction of neutrons and protons in overlapping orbitals [6]. Shell model calculations with realistic forces are succesful in reproducing the measured binding energies [7], whereas with Skyrme forces, no Wigner term appears in Hartre-Fock-Bogolyubov calculations and only a small one in Hartree-Fock calculations [8]. However, by invoking a particular isoscalar pairing force that

breaks geometric symmetries, Satula and Wyss obtain a significant Wigner energy in approximately number-projected Bogolyubov calculations [9]. In finite-temperature Bogolyubov calculations with a Yamaguchi force, Röpke *et al.* get at low temperature in the local density approximation a contribution from neutron-proton pairing to the binding energies in the $A = 40$ isobaric chain with a maximum at $N - Z = -1$ and a quite irregular dependence on the asymmetry [10].

The RPA is the leading order correction to the Hartree-Fock-Bogolyubov approximation. I have therefore calculated the RPA correlation energy from the schematic Hamiltonian

$$H = H_0 - G\mathbf{P}^\dagger \cdot \mathbf{P} + \frac{\kappa}{2}\mathbf{T}^2, \quad H_0 = \sum_{k\sigma\tau} \epsilon_k a_{k\sigma\tau}^\dagger a_{k\sigma\tau}.$$

In this expression, the index $k\sigma\tau$ labels orthonormal nucleon orbitals, and $a_{k\sigma\tau}$ are the corresponding annihilation operators. $k\sigma$ takes the values k and $\bar{k}$ so that the orbital $\bar{k}$ is obtained from the orbital k by time-reversal, and τ is ‘n’ for a neutron orbital or ‘p’ for a proton orbital. $\mathbf{P}$ is the pair annihilation isovector, and $\mathbf{T}$ denotes the total isospin. The former has the coordinates

$$P_x = (-P_n + P_p)/\sqrt{2}, \quad P_y = -i(P_n + P_p)/\sqrt{2}, \quad P_z = P_{np},$$

$$P_\tau = \sum_k a_{\bar{k}\tau} a_{k\tau}, \quad P_{np} = \sum_k (a_{\bar{k}p} a_{kn} + a_{\bar{k}n} a_{kp})/\sqrt{2}.$$

The single-nucleon energy ϵ_k takes Ω equidistant values separated by η , and G and κ are coupling constants.

To describe states with a given number $A_v = \sum_{k\sigma\tau} a_{k\sigma\tau}^\dagger a_{k\sigma\tau}$ of valence nucleons and a given isospin, I employ the Routhian

$$R = H - \lambda A_v - \mu T_z.$$

It is just for convenience that T_z is chosen here as the isospin-coordinate to be constrained. Since H is isobarically invariant, one could equivalently constrain the projection of $\mathbf{T}$ on any axis in isospace. Following Marshalek [11] I base the RPA on the Hartree-Bogolyubov (not Fock) self-consistent state derived from R . This is the Bogolyubov vacuum that minimizes $E_0 - \lambda\langle A_v \rangle - \mu\langle T_z \rangle$, where

$$E_0 = \langle H_0 \rangle - \frac{|\Delta|^2}{G} + \frac{\kappa}{2}\langle \mathbf{T} \rangle^2, \quad \Delta = -G\langle \mathbf{P} \rangle.$$

At the minimum one has $\langle T_x \rangle = \langle T_y \rangle = 0$. For large values of μ a product of neutron and proton BCS states is expected. Since this state is invariant under

the transformation $\exp(-i\pi(A_v/2+T_z))$, I enforce this symmetry, which entails $\Delta_z = 0$. I furthermore assume that both gaps $\Delta_n = (-\Delta_x + i\Delta_y)/\sqrt{2}$ and $\Delta_p = (\Delta_x + i\Delta_y)/\sqrt{2}$ are positive, as may always be achieved by a transformation of the form $\exp(-i(\xi A_v + \chi T_z))$. When λ is placed midway between the lowest and the highest ϵ_k , one then gets $\langle A_v \rangle = 2\Omega$ and $\Delta_n = \Delta_p \equiv \Delta$ for any value of μ due to the equidistant single-nucleon spectrum. To speed up the calculation, I keep Δ rather than G fixed with the variation of μ . Then G varies in the case considered by less than .6 %.

With quasinucleon annihilation operators $\alpha_i = \sum_{k\sigma\tau} (u_{i,k\sigma\tau} a_{k\sigma\tau} + v_{i,k\sigma\tau} a_{k\sigma\tau}^\dagger)$ defined by

$$[\alpha_i, R_m] = E_i \alpha_i, \quad E_i > 0, \quad \{\alpha_i, \alpha_j^\dagger\} = \delta_{ij},$$

$$R_m = H_0 + \Delta(P_n + P_n^\dagger + P_p + P_p^\dagger) + \kappa \langle T_z \rangle T_z - \lambda A_v - \mu T_z,$$

the RPA Routhian $\tilde{R}$ is obtained by truncating to second order the boson expansion of R that results from making in the expressions for H_0 , $\mathbf{P}$, $\mathbf{T}$ and A_v the substitutions

$$\alpha_j \alpha_i = b_{ij} + \dots, \quad \alpha_i^\dagger \alpha_j = \sum_l b_{il}^\dagger b_{jl},$$

where the boson annihilation operators $b_{ij} = -b_{ji}$ satisfy $[b_{ij}, b_{lm}] = 0$ and $[b_{ij}, b_{lm}^\dagger] = \delta_{il}\delta_{jm} - \delta_{im}\delta_{jl}$. The normal mode annihilation operators $B_\nu = \sum_{i<j} (\phi_{\nu,ij} b_{ij} + \psi_{\nu,ij} b_{ij}^\dagger)$ are then given by

$$[B_\nu, \tilde{R}] = \omega_\nu B_\nu, \quad \omega_\nu > 0, \quad [B_\nu, B_\rho^\dagger] = \delta_{\nu\rho},$$

and the ground state energy is [11] $E = E_0 + E_2$ with

$$E_2 = \sum_{k\tau\tau'} \left(-G |\langle \alpha_{\bar{k}\tau'} \alpha_{k\tau} \mathbf{P} \rangle|^2 + \frac{\kappa}{2} |\langle \alpha_{\bar{k}\tau'} \alpha_{k\tau} \mathbf{T} \rangle|^2 \right) + \frac{1}{2} \left(\sum_\nu \omega_\nu - \sum_{i<j} [b_{ij}, [\tilde{R}, b_{ij}^\dagger]] \right).$$

It may be noticed that when the terms with four quasinucleon annihilation operators or four quasinucleon creation operators are removed from the expression for the Hamiltonian H so that the Bogolyubov vacuum becomes an eigenstate of H (when the operators A_v and T_z are replaced by their eigenvalues in the relation $H = R + \lambda A_v + \mu T_z$), then only the first term in the expression for E_2 survives, and $E_0 + E_2$ becomes an exact expression for the




Fig. 1. $E_0 - E_{0,T=0}$, $E_2 - E_{2,T=0}$ and $\mu/2$ as functions of T . $E_0 - E_{0,T=0}$ is scaled by the factor $1/3$.

ground state energy. Therefore this expression derived from the boson expansion is expected to provide a reliable approximation even though most of the normal modes are not much different from two-quasinucleon excitations.

The implications of the symmetries of this model for the normal modes are discussed by Ginocchio and Wesener [12]. Two Goldstone modes result from the commutation relations $[A_v, R] = [T_z, R] = 0$. Furthermore, since $[T_+, R] = \mu T_+$ one normal mode has the frequency μ . Its annihilation operator is the linear boson part of $T_+/\sqrt{2\langle T_z \rangle}$, and it becomes a Goldstone mode in the limit $\mu \rightarrow 0$. The degree of freedom of this mode is the direction of the isospin. In particular the isospin quantum numbers $M_T = T = \langle T_z \rangle$ may be assigned to the ground state of the Routhian R . The Goldstone modes contribute with the frequency zero to the expression for the second order energy E_2 .

The parameters of the calculation are chosen so as to simulate the $A = 48$ isobaric chain: $\Omega = 24$, $\eta = 2.1$ MeV, $\Delta = 1.7$ MeV, $\kappa = 1.2$ MeV. The result is shown in Figure 1. $E_0 - E_{0,T=0}$ depends essentially quadratically on T . It is in fact given in a very good approximation by the expression $E_0 - E_{0,T=0} = \frac{1}{2}(\eta + \kappa)T^2$ obtained for $\Delta = 0$. The almost exactly quadratic T -dependence of $E_0 - E_{0,T=0}$ is seen also indirectly from the linearity of $\mu = dE_0/dT + 2(\Delta/G)^2 dG/dT$, where the second term is negligible. $E_2 - E_{2,T=0}$ shows a different behaviour. It rises for $T \approx 0$ linearly with T and is in fact in this limit equal to $\mu/2$. The linearity thus stems from the single term in the expression for E_2 which represents the zero-point energy of the normal mode with the frequency μ , or, in other words, from the quantal fluctuation of the

isospin.

The rest of the second order energy E_2 is fairly independent of T . This suggests that the other normal modes are to a large extent independent of the iso-rotational degree of freedom, or, stated otherwise, that the iso-rotation is highly collective. The deformation underlying this collectivity is in the pair field [13]. Thus with $\Delta_z = 0$ the isovector Δ is perpendicular to the iso-rotational axis. So it breaks the iso-rotational invariance with respect to this axis, and a collective iso-rotation can arise. On the other hand the contribution to E_2 from the non-collective modes varies from $T = 0$ to $T = 4$ by almost 2 MeV, so the RPA correlation energy should be taken into account in a detailed comparison of the results of Hartree-Fock-Bogolyubov calculations with the empirical masses.

With $E_0 - E_{0,T=0} = \frac{1}{2}(\eta + \kappa)T^2$ and $E_2 - E_{2,T=0} = \frac{1}{2}\mu = \frac{1}{2}dE_0/dT = \frac{1}{2}(\eta + \kappa)T$ we have altogether $E - E_{T=0} = \frac{1}{2}(\eta + \kappa)T(T + 1)$, that is, we get the T -dependence of the symmetry energy found in the data. The Hartree-Fock-Bogolyubov energy expectation value includes only the first sum in the expression for E_2 [11]. This sum is cancelled to a large extent by parts of the second term in the second sum. For the contributions from the symmetry potential energy $\frac{\kappa}{2}T^2$ this cancellation is in fact exact. For $\Delta > 0$ the first sum in the expression for E_2 is as a function of T analytic and even at $T = 0$. For $\Delta = 0$ it equals, however, $\frac{\kappa}{2}T$. Thus it produces in the absence of pairing a Wigner term, albeit with only $\kappa/(\eta + \kappa) \approx 35\%$ of the full value. Although the forces there are different, this may explain the experience with Hartree-Fock-Bogolyubov and Hartree-Fock calculations mentioned in the introduction.

A Wigner term corresponding to the term $\frac{\kappa}{2}T$ in the present model is actually the only one that may be derived from arguments like those in References 2–4, which are based on the form of the residual two-nucleon interaction. It is remarkable that with a deformation one gets also a term $\frac{\eta}{2}T$ corresponding to the ‘kinetic’ part $\frac{\eta}{2}T^2$ of the symmetry energy.

The symmetry potential energy $\frac{\kappa}{2}T^2$ of the present model differs from an interaction potential energy $\kappa \sum_{i < j} \mathbf{t}_i \cdot \mathbf{t}_j$, where the index i or j labels the individual nucleons, by the term $\frac{3}{8}\kappa A_v$, which depends only on the number A_v of valence nucleons. This interaction favours isoscalar nucleon pairs. It is included in the Hamiltonian in order to get a realistic symmetry energy coefficient. It is in fact well known and discussed in detail by for example Bohr and Mottelson [4] that the kinetic term accounts for only a part of the empirical symmetry energy coefficient. Bohr and Mottelson consider a contribution to the single-nucleon potential energy corresponding to a two-nucleon potential of the form $\kappa \mathbf{t}_1 \cdot \mathbf{t}_2$. Thus the form of the symmetry potential energy of the present model is adopted from their discussion. This symmetry force is inessential, however, for the formation of the Wigner energy, which

results, as it was seen, from the collectivity of the iso-rotation due to the isobaric non-invariance of the isovector pair field.

I am indebted to Stefan Frauendorf for drawing my attention to the issue of the Wigner energy and for discussions of the matter.

References

- [1] W.D. Myers, W.J. Swiatecki, Nucl. Phys. A 81 (1966) 1.
- [2] E.P. Wigner, Phys. Rev. 51 (1937) 106.
- [3] I. Talmi, Rev. Mod. Phys. 34 (1962) 704.
- [4] A. Bohr, B.R. Mottelson, Nuclear Structure, Vol. 1, Benjamin, New York, 1969
- [5] J.Duflo, A.P. Zuker, Phys. Rev. C 52 (1995) R23.
- [6] W.D. Myers, W.J. Swiatecki, Nucl. Phys. A 612 (1997) 249.
- [7] E. Caurier, G. Martinez-Pinedo, F. Nowacki, A.Poves, J. Retamosa, A.P. Zuker, Phys. Rev. C 59 (1999) 2033.
- [8] W. Satula, R.A. Wyss, Acta Phys. Polon. B 32 (2001) 2441.
- [9] W. Satula, R.A. Wyss, Phys. Lett. B 393 (1997) 1.
- [10] G. Röpke, A. Schnell, P. Schuck, U. Lombardo, Phys. Rev. C 61 (2000) 024306.
- [11] E.R. Marshalek, Nucl. Phys. A 275 (1977) 416.
- [12] J.N. Ginocchio, J. Wesener, Phys. Rev. 170 (1968) 859.
- [13] S.G. Frauendorf, J.A. Sheikh, Nucl. Phys. A 645 (1999) 509.